



For Back Up Use



This copy of the Orderset is for Information & BackUp purposes only.

It is intended for use during downtimes.

**For daily use, please access the version on
EntryPoint**

To launch EntryPoint, go to PCI (Patient Care Inquiry) in Meditech. Look for “EntryPoint Launch” and follow the instructions. Log in using your PACS (or Outlook) username and password in lowercase.

Thanks.



Height _____ cm Weight _____ kg

Allergies _____

Adult Emergency Department (ED) Syphilis Order Set		M	K	O
Orders Processed Date (dd/mm/yyyy)	Consults			
	<input type="checkbox"/> Infectious Disease: For high risk patients (e.g. pregnant patients, immunocompromised, receiving HIV Prophylaxis)			
Time (hhmm)	Lab Investigations			
	As Indicated (females, age 10 – 55)			
	<input type="checkbox"/> β HCG			
By	If not already done			
	<input checked="" type="checkbox"/> CBC <input checked="" type="checkbox"/> Creatinine <input checked="" type="checkbox"/> ALT, ALP, Bilirubin			
	<input checked="" type="checkbox"/> HCV Serology <input checked="" type="checkbox"/> HIV Serology <input checked="" type="checkbox"/> VDRL (Syphilis) Serology			
	<input checked="" type="checkbox"/> Hep B Surface Antibody, Hep B Surface Antigen			
Status	Lab Investigations below to be done Prior to First Dose of Antibiotic			
	<input type="checkbox"/> Urine sample (first void) for nucleic acid amplification test (NAAT) testing for Gonorrhea and Chlamydia (Both males and females)			
	<input type="checkbox"/> Vaginal swab/endo-cervical NAAT testing for Gonorrhea and Chlamydia			
Processing Reviewed by	<input type="checkbox"/> Rectal/pharyngeal NAAT testing for Gonorrhea and Chlamydia			
	<input type="checkbox"/> Urethral Culture for Gonorrhea and Chlamydia (Both male or female)			
	<input type="checkbox"/> Endo-cervical, vaginal, rectal, or pharyngeal culture for Gonorrhea and Chlamydia			
Status	<input type="checkbox"/> Genital swab for HSV1, HSV2 and Varicella PCR			
	<input type="checkbox"/> Genital/non-genital swab for Chlamydia culture			
	<input type="checkbox"/> Other: _____			
Faxed by	For Confirmed Syphilis			
	** Penicillin G Benzathine is the only effective drug for the treatment of syphilis during pregnancy **			
	Ensure the correct Penicillin G Benzathine is used			
	<input type="checkbox"/> penicillin G benzathine 2.4 million units (2 syringes of 1.2 million units) intramuscularly x 1 dose			
	<input type="checkbox"/> Inject one 1.2 million unit syringe into each ventrogluteal site			
	Documented Allergy to Penicillin			
	<input type="checkbox"/> doxycycline 100 mg PO BID for 14 days			
	Additional Orders			

Telephone Order _____
 Ordering Practitioner, Designation Signature Date/Time (dd/mm/yyyy hhmm)

If Telephone Order _____
 Ordering Physician Date (dd/mm/yyyy) Time (hhmm) Read Back



Chart Copy – Do Not Destroy

Rev. 09/2022/V1 ORD338